

Effect of crop residue management on N₂O emissions in European cropping systems

Carozzi M.¹, Haas E.², Scheer C.², Butterbach-Bahl K.², Loubet B.¹, Massad R.S.¹

¹ EcoSys, INRA, Université Paris Saclay, Thiverval-Grignon, France

² IMK-IFU, Karlsruhe Institute of Technology, Garmisch-Partenkirchen, Germany

E-mail: marco.carozzi@inra.fr

Abstract

Crop residue handling can significantly affect N₂O emissions. Four different scenarios of crop residues management were evaluated with two biogeochemical models in the European cropping systems. Preliminary results show the positive effect of nitrogen addition by residues on crop yields, highlighting a potential mitigation of N₂O emissions accompanied with rational fertilisation.

Crop residue management, nitrous oxide, cropping systems, ecosystem modelling

1. Introduction

Crop residue management is crucial to maintain or improve soil organic carbon and fertility. Their handling on the field can significantly affect nitrous oxide (N₂O) production. Ecosystem models represent a suitable tool to simulate biogeochemical processes in both time and space, facilitating the analysis of management scenarios. The aim of this work is to reproduce N₂O emissions from different management of crop residue in the European cropping systems.

2. Materials and methods

Two ecosystem models, Landscape-DNDC (Haas et al., 2012) and CERES-EGC (Gabrielle et al., 1995), were applied in the EU-27 cropping systems from 1970 to 2010. Four residue managements were evaluated: i) exported, ii) left on the surface, iii) incorporated into the soil, iv) exported or incorporated in function of the crop species.

A spatialized dataset (0.25° × 0.25° latitude-longitude simulation units) composed by climate, soil and crop data was created to run the models. Daily climate data were provided by Sheffield et al., (2006), whereas soil characteristics were extracted from the European Soil Database. Crop species and management data were obtained from Eurostat statistics, retaining the two most frequent rotations for each simulation unit. N fertilisation ranged from

a few dozen to 370 kg N ha⁻¹ y⁻¹ and was applied to 91% of the EU-cropping systems. Crops were automatically irrigated.

3. Results and discussion

Preliminary results with CERES-EGC model show a reduction of N₂O emissions of about 30% when the residues were exported from the field (i) compared to the residue management in function of the crop species (iv). Residue incorporation (iii) marks a slight increase of N₂O compared to the scenario (iv). Crop yields were 13% lower and 15% higher for the exported and incorporated scenarios, respectively compared with scenario (iv).

4. Conclusions

These results underline the effects of N addition with residue incorporation on yields, highlighting a potential mitigation of N₂O emissions with rational fertilisation.

Acknowledgements

This work was carried out in the framework of the project ResidueGAS (FACCE ERA-GAS)

References

- Gabrielle B., Menasseri S., Houot S., 1995. Analysis and field- evaluation of the CERES models' water balance component. *Soil Sci Soc Am J* 59:1402–1411.
- Haas E., Klatt S., Fröhlich A., Kraft P., Werner C., Kiese R., Grote R., Breuer L., Butterbach-Bahl K., 2012. Landscape-DNDC: a process model for simulation of biosphere–atmosphere–hydrosphere exchange processes at site and regional scale. *Landscape Ecol.*, 28, 615–636.
- Sheffield, J., G. Goteti, and E. F. Wood. 2006. Development of a 50-yr high-resolution global dataset of meteorological forcings for land surface modeling, *J. Climate*, 19 (13), 3088-3111.